

**Stop taking the following medications at least 2 weeks before surgery due to the fact these drugs can increase the risk of bleeding during and after surgery. Continue other medications as directed by your physician.**

**\*\*PLEASE CHECK WITH YOUR PRIMARY PHYSICIAN  
BEFORE STOPPING THESE MEDICATIONS\*\***

<b>Advil</b>	Bufferin	<b>Eliquis</b>	Midol	<b>Orudis</b>	Xarelto
Aggrenox	<b>Cangrelor</b>	Feldene	<b>Midrin</b>	Pediaprofen	
<b>Aleve</b>	Celebrex	<b>Fish Oil</b>	Mobic	<b>Plavix</b>	
Alka-Seltzer	<b>Clinoril</b>	Goody's Powder	<b>Motrin</b>	Relafen	
<b>Anaprox</b>	Coumadin	<b>Ibuprofen</b>	Nalfen	<b>Soma Compound</b>	
Aspergun	<b>Daypro</b>	Indocin	<b>Naprelan</b>	Stan Back	
<b>Aspirin</b>	Dolobid	<b>Lodine</b>	Naprosyn	<b>Torodol</b>	
B C Powder	<b>Effient</b>	Meloxicam	<b>Nuprin</b>	Voltaren	

**You CAN take Tylenol, Lortab or Codeine. If you have a question as to if a medication contains aspirin, please contact your pharmacist.**

**Generally, we recommend that you take a supplemental Vitamin C, Iron or a Multi-Vitamin twice a day to aid in the healing process.**